



Share on Facebook



Share on Twitter



Share via email


[About](#) [Our Bloggers](#) [Buy Something](#)
[Subscribe by Email](#) [Subscribe to RSS](#) 

## Sandia and UNM lead effort to destroy cancers

on APRIL 18, 2011

### [Non-Hodgkin's Lymphoma](#)

Have you been recently diagnosed?  
 Learn more about the BRIGHT study.  
[www.treandaclinicalresearch.com](http://www.treandaclinicalresearch.com)

### [HTD Biosystems Inc.](#)

Formulation Development of Proteins  
 Liposomes & Freeze-dried Products  
[www.htdcorp.com](http://www.htdcorp.com)

### [Lung cancer?](#)

Compensation trust fund information  
 Find out if you qualify  
[www.calldavid.com](http://www.calldavid.com)



Ads by Google

ALBUQUERQUE, N.M. — Melding nanotechnology and medical research, Sandia National Laboratories, the University of New Mexico, and the UNM Cancer Research and [Treatment Center](#) have produced an effective strategy that uses nanoparticles to blast cancerous cells with a mélange of killer drugs.

In the cover article of the May issue of *Nature Materials*, available online April 17, the [researchers](#) describe silica nanoparticles about 150 nanometers in diameter as honeycombed with cavities that can store large amounts and varieties of drugs.

“The enormous capacity of the nanoporous core, with its high surface area, combined with the improved targeting of an encapsulating lipid bilayer [called a liposome], permit a single ‘protocell’ loaded with a drug cocktail to kill a drug-resistant [cancer cell](#),” says Sandia researcher and UNM professor Jeff Brinker, the principal investigator. “That’s a millionfold increase in efficiency over comparable methods employing liposomes alone — without nanoparticles — as drug carriers.”

The nanoparticles and the surrounding cell-like membranes formed from liposomes together become the combination referred to as a protocell: the membrane seals in the deadly cargo and is modified with molecules (peptides) that bind specifically to receptors overexpressed on the cancer cell’s surface. (Too many receptors is one signal the cell is cancerous.) The nanoparticles provide stability to the supported membrane and contain and

 Google™ Custom Search

### FROM OUR BLOGGERS

- [What do extreme weather events tell us about climate change?](#)
- [Track Kepler's Candidate Exoplanets with a new app](#)
- [Out of this world children's science books](#)
- [DNA Day and Body Modification](#)
- [For Science History Buffs: Blood Work by Holly Tucker](#)

[Buy a Blogad!](#)

blog advertising is good for you

release the therapeutic cargo within the cell.

A current Food and Drug Administration-approved nanoparticle delivery strategy is to use liposomes themselves to contain and deliver the cargo. In a head-to-head comparison of targeted liposomes and protocells with identical membrane and peptide compositions, Brinker and colleagues [report](#) that the greater cargo capacity, stability and targeting efficacy of protocells leads to many times greater cytotoxicity [destruction] directed specifically toward human [liver cancer](#) cells.

Another advantage to protocells over liposomes alone, says lead author Carlee Ashley, a Harry S. Truman post-doctoral fellow at Sandia's [California](#) site in Livermore, is that liposomes used as carriers need specialized loading strategies that make the process more difficult. "We've demonstrated we can just soak nanoparticles to load them with unique drug combinations needed for personalized medicine. They effectively encapsulate toxins as well as siRNA [ribonucleic acid] that silence expressions of proteins."

RNA, the biological messenger that tells cells which proteins to manufacture, in this case is used to silence the cellular factory, a way of causing apoptosis or cell death. "Si" is short for "silence."

The lipids also serve as a shield that restricts toxic [chemotherapy](#) drugs from leaking from the nanoparticle until the protocell binds to and takes hold within the cancer cell. This means that few poisons leak into the system of the human host, if the protocells find no cancer cells. This cloaking mitigates toxic [side effects](#) expected from conventional chemotherapy.

Instead, the particles — crafted small enough to float under the radar of the liver and other cleansing organs — can circulate harmlessly for days or weeks, depending on their engineered size, seeking their prey.

A library of phages — [viruses](#) that attack bacteria — was created at UNM's nationally accredited cancer center by collaborator David Peabody. This permitted researchers to expose the phages to a group of cancerous cells and normal cells, allowing [identification](#) of peptides that bind specifically to cancer cells but not normal cells.



You like this. · Admin Page · Insights · Error You like this Sign Up to

"Proteins modified with a targeting peptide that binds to a particular carcinoma exhibit a

10,000-fold greater affinity for that cancer than for other unrelated cells," Ashley said.

Brinker adds, "A key feature of our protocell is that its fluid bilayer allows high-affinity binding with just a few of these peptides

overall. This reduces nonspecific binding and immune response.”

The method is being tested on human cancer cells in vivo, and will shortly be tested on mouse tumors at UNM’s cancer center.

The researchers continue to optimize the size of the porous silica particle, which is formed by aerosolizing a precursor solution. The porous nanoparticle fabrication process — called evaporation-induced self-assembly, and pioneered in the Brinker lab — produces particles from 50 nm to several microns in diameter. Particle [sizes](#) between 50 and 150 nanometers in diameter are ideal for maximizing circulation and uptake into cancer cells, so the particles are preselected by size before their formation into protocells.

“Their overall dimensions determine how widely they’ll be distributed in the bloodstream,” Brinker said. “We’re altering our synthesis to favor the smaller sizes.”

Also of importance to the circulation time of the particle are its electrical charge and hydrophobicity [avoidance of water], which can improve or detract from its ability to remain free of unwanted molecular or energetic entanglements.

The method may be commercially available in five years, researchers estimate.

Brinker is a Sandia Fellow and UNM Regent’s and Distinguished Professor of Chemical and Nuclear Engineering and member of UNM’s cancer center.

Other institutions involved in the research include the University of California Davis, and the University of Waterloo in Canada.

Funding was provided by the [National Cancer Institute](#), the National Science Foundation, the Department of Energy’s Basic Energy Sciences program, the Air Force Office of Scientific Research, and Sandia’s Laboratory Directed Research and Development office.

The work is the first to show targeted delivery of nanoparticles to cancers that is supported in part by a grant from NCI’s Alliance for Nanotechnology in Cancer.

#### [Metastatic tumors](#)

New Mexico’s Gamma Knife Center for treatment of metastatic tumors.

[www.lovelace.com](http://www.lovelace.com)

#### [Non-Hodgkin's Lymphoma](#)

Have you been recently diagnosed? Learn more about the BRIGHT study.

[www.treandaclinicalresearch.com](http://www.treandaclinicalresearch.com)

#### [Lymphoma Treatments](#)

Learn about innovative treatments for

Lymphoma diagnosis today.  
[CancerCenter.com/CareThatNeverQuits](http://CancerCenter.com/CareThatNeverQuits)



Ads by Google

## Subscribe

Subscribe to our e-mail newsletter to receive updates.



← 'I'm a Mac' — so what? Study finds way to measure brand personality appeal  
Researchers pinpoint graphene's varying conductivity levels →

No comments yet.

## Leave a Reply

Name (required)

Mail (will not be published) (Required)

Website



CAPTCHA Code \*

[Much Older Archives](#)

